Theodore B. Reinhart, P.E.

- 2011-now: Energy-PE, LLC
- 1985-2011: Laclede Gas Company
- 1984-1985: Londe-Parker-Michels
- M.S. 1984: University of Missouri – Columbia,
  Mechanical and Aerospace Engineering
- 1981-1982 & 1983: Industrial Engineering and Equipment Co.
- 1979-1981: Mary Institute
- B.A. 1978: Rice University, Physics

energy-pe.com @energy_pe



The CO_2 Answer in 6 Easy Steps

A presentation based on the RealClimate post

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Learning Objectives

- Explain the difference between predictions and data
- Describe how greenhouse gases affect surface temperature and the effect of changing the greenhouse gas effectiveness (1)
- Understand how spectroscopy measures the radiative transfer absorbed by trace greenhouse gases (2)
- Demonstrate the context of industrial age concentrations of trace greenhouse gases (3)
- Use radiative forcing and climate sensitivity to calculate how the environment reacts to change (4, 5, 6)

asu.cas.cz/~bezdek/vyzkum/rotating_3d_globe/

earthobservatory.nasa.gov/Features/Milankovitch/milankovitch_3.php



Astro Bob



26 000 & 40 000 years — NASA — 100 000 years — USC

The obliquity and eccentricity of the Earth

Aphelion 152 098 232 km (first week of July, summer in northern hemisphere)

Perihelion 147 098 290 km (first week of January, winter in northern hemisphere)

The Earth is about 5 billion meters closer to the Sun at perihelion than at aphelion

$$\frac{A^2}{P^2} = 1.069$$



scienceblogs.com

Disk and sphere of same radius catch the same solar radiation

Area of disk = πr^2

Area of sphere = $4\pi r^2$

Ratio = 1:4

albedo: The fraction of power scattered back out into space from the total radiation incident on an astronomical body.

globalwarmingart.com/wiki/File:65_Myr_Climate_Change_Rev_png



Cenozoic (new life) "Age of mammals" **Paleocene, Eocene, Oligocene, Miocene, Pliocene, Pleistocene, and Holocene (not labeled)**

Less than 1.5% of the age of the Earth

60 mya is oldest Powder River Basin coal



Wikimedia

Lake Vostok ice core



derivative work: Alexchris Wikimedia

J. R. Petit, et al, "Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica" (1999)

nature.com/nature/journal/v399/n6735/fig_tab/399429a0_F3.html



derivative work: Autopilot Wikimedia

J. R. Petit, et al, "Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica" (1999)

http://www.nature.com/nature/journal/v399/n6735/fig_tab/399429a0_F2.html

Quiz: Heat Transfer (Part One)

For **each of these three** mechanisms:

- **Conduction**
- **Convection**
- **Radiation**



What percentage is it of the total heat transfer

from the Sun to the Earth?



eesc.columbia.edu

Solar "constant," mean total solar irradiance,
***TSI* = 1366 W/m^2 (1.36 x 10^6 erg/cm^2 sec)**

Instantaneous insolation cycles daily, annually;

obliquity, eccentricity, changes cycle long term.

Quiz: Heat Transfer (Part Two)

- **Conduction**
- **Convection**
- **Radiation**

Which one of these three mechanisms is **100%** of the total heat transfer

from the Earth to space?

The CO_2 Answer in 6 Easy Steps

1. **There is a natural greenhouse effect**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Mean surface temperature, $\boldsymbol{T_s}$, **~15°C**

produces upward surface flux of longwave radiation

$$\boldsymbol{G = \sigma T_s^4 \cong 390 \frac{W}{m^2}}$$

Stefan-Boltzmann constant:

$\sigma = \frac{2\pi^5 k^4}{15 c^2 h^3} = 5.67 \text{ x } 10^{-8}\ W m^{-2} K^{-4}$

Net solar radiation absorbed

$$\boldsymbol{S = \frac{(1 - a) TSI}{4} \cong 240 \frac{W}{m^2}}$$

$a = 0.306$

$TSI = 1366\ W/m^2$

means 240 W/m^2 of longwave radiation is emitted

Atmosphere must absorb **150 $\boldsymbol{W/m^2}$** net

a number that would be zero without greenhouse gases

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Tidal power
3.7 TW
0.007 W/m^2

Geothermal power
44.2 TW
0.087 W/m^2

Wind power
72 TW
0.14 W/m^2



Solar power
121,000 TW
237 W/m^2

World power use
17.5 TW
0.034 W/m^2

http://www.smu.edu/~/media/Site/Dedman/Academics/Programs/Geothermal%20Lab/Graphics/SMUHeatFlowMap2011_CopyrightVA0001377160_jpg.ashx?la=en

ces.fau.edu/nasa/module-2/atmosphere/earth.php







Layers in the Atmosphere



realclimate.org

$$S = (1 - a)TSI/4$$

$$G = \sigma T_s^4$$

$$\lambda A = \lambda \sigma T_a^4$$

λ = **emissivity**, effectively the strength of the greenhouse effect

Equations of equilibrium

Surface: $S + \lambda A = G$

Atmosphere: $\lambda G = 2\lambda A$

Planet: $S = \lambda A + (1 - \lambda)G$

$$G = \sigma T_s^4$$

$$= \frac{S}{1 - 0.5\lambda}$$

$$\lambda = 0.769$$

$$T_s = 288°K$$

$$T_s = 15°C$$



realclimate.org

$$A = \sigma T_a^4 = \frac{S}{2 - \lambda}$$

$$T_a = 242°K$$

$$T_a = -31°C$$

λ	G	A	T_s	T_a
0	240 W/m^2	-	$-18°C$	-
0.667	360 W/m^2	180 W/m^2	$9°C$	$-36°C$
0.769	390 W/m^2	195 W/m^2	$15°C$	$-31°C$
1	480 W/m^2	240 W/m^2	$30°C$	$-18°C$





NASA

science-edu.larc.nasa.gov/energy_budget/

The CO_2 Answer in 6 Easy Steps

1. **There is a natural greenhouse effect**



The CO_2 Answer in 6 Easy Steps

2. **Trace gases contribute to the natural greenhouse effect**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/



royalsocietypublishing.org

William Hyde Wollaston, A Method of Examining Refractive and Dispersive Powers, by Prismatic Reflection (1802)

Day-light - dark room - crevice 1/20" - distance 10’ or 12’- through a prism of flint-glass, free from veins held near the eye - beam is separated into four colours only - red, yellowish green, blue, and violet; in the proportions in Fig. 3.

Other distinct dark lines, f and g, might be mistaken for the boundary of these colours. [bottom of page 378]

rstl.royalsocietypublishing.org/content/92/365.full.pdf





astro.umontreal.ca/~paulchar/grps/histoire/newsite/sp/great_moments_e.html

Wikipedia



Fraunhofer lines (1814)



planetaryvision.blogspot.com/2013/01/the-fallacy-of-greenhouse-effect-4.html Planetary Vision



Ångström's spectrometer (1868)

iris.univ-lille1.fr/bitstream/handle/1908/1400/Q11406_1.pdf

1849 - Jean Bernard Léon Foucault
1853 - Anders Jonas Ångström
Showed the emission and absorption lines matched for a given material

Wikipedia, Jan Homann

$$\lambda = \text{const.} \left(\frac{m^2}{m^2 - n^2} \right)$$



Wikipedia, JabberWok

1853 - Ångström discovered four visible lines of hydrogen

410, 434, 486, and 656 nm

The Balmer series

1885 - Johann Jacob Balmer discovered relationship between wavelengths; predicted others

1913 - Niels Henrik David Bohr quantum theory of the atom



An optical disc in fluorescent light

The major Fraunhofer lines and their elements

Designation	Element	Wavelength (nm)	Designation	Element	Wavelength (nm)
y	O_2	898.765	c	Fe	495.761
Z	O_2	822.696	F	Hβ	486.134
A	O_2	759.370	d	Fe	466.814
B	O_2	686.719	e	Fe	438.355
C	Hα	656.281	G'	H	434.047
a	O_2	627.661	G	Fe	430.790
D_1	Na	589.592	G	Ca	430.774
D_2	Na	588.995	h	Hδ	410.175
D_3 or d	He	587.5618	H	Ca^+	396.847
e	Hg	546.073	K	Ca^+	393.368
E_2	Fe	527.039	L	Fe	382.044
b_1	Mg	518.362	N	Fe	358.121
b_2	Mg	517.270	P	Ti^+	336.112
b_3	Fe	516.891	T	Fe	302.108
b_4	Mg	516.733	t	Ni	299.444

Wikipedia

1824 Jean-Baptiste Joseph Fourier

Step 1. There is a natural greenhouse effect

1859 John Tyndall

Step 2. Trace gases contribute to the natural greenhouse effect



1879 Josef Stefan

On the basis of experimental measurements made by John Tyndall

1884 Ludwig Boltzmann

Derived from theoretical considerations, using thermodynamics

1896 Svante Arrhenius

Burning fossil fuels increases atmospheric CO2 and the greenhouse effect

http://www.skepticalscience.com/history-climate-science.html
https://www.aip.org/history/climate/summary.htm

planetaryvision.blogspot.com/2013/01/the-fallacy-of-greenhouse-effect-4.html



From: www.barrettbellamyclimate.com/page15.htm

"… using line-by-line radiative transfer codes … removing the effect of CO_2 reduces the net LW absorbed by **~14%, or** around **30 W/m^2**."

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/



Geostationary Operational Environmental Satellite - 1

Launched: October 16, 1975 Deactivated: March 7, 1985

cimss.ssec.wisc.edu/goes/comet/radiative_transfer.html

goes.gsfc.nasa.gov/text/history/goes/goes.html

The CO_2 Answer in 6 Easy Steps

2. Trace gases contribute to the natural greenhouse effect





http://serc.carleton.edu/quantskills/methods/quantlit/BigNumbers.html

The CO_2 Answer in 6 Easy Steps

3. **The trace greenhouse gases have increased markedly due to human emissions**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

"CO_2 is up more than 30%,
CH_4 has more than doubled,
N_2O is up 15%,
tropospheric O_3 has also increased.
New compounds such as halocarbons (CFCs, HFCs) did not exist in the pre-industrial atmosphere."

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/



NOAA

youtube.com/watch?v=vA7tfz3k_9A

Youtube movie of a time history of atmospheric carbon dioxide from 800,000 years ago until January, 2012, Earth System Research Laboratory, NOAA

http://www.esrl.noaa.gov/gmd/ccgg/trends/history.html

The CO_2 Answer in 6 Easy Steps

3. **The trace greenhouse gases have increased markedly due to human emissions**



The CO_2 Answer in 6 Easy Steps

4. **Radiative forcing is a useful diagnostic and can easily be calculated**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

If the sun became stronger by 2%

TOA radiation balance would change by

$$\mathbf{0.02 \times 1366 \times 0.7/4 = 4.8\, W/m^2}$$

(taking albedo and geometry into account)

This is the **radiative forcing** (RF). Changes in greenhouse absorbers, albedo have analogous impacts



$$\boldsymbol{S = \frac{(1-a)TSI}{4}}$$

$$\boldsymbol{\cong 245\ W/m^2}$$

$$\boldsymbol{T_s = \sqrt[4]{\frac{(1-a)TSI}{4\sigma(1-0.5\lambda)}}}$$

$$\boldsymbol{T_s \cong 289.5°K = 16.5°C}$$

$a = 0.306$ $TSI = 1366\ W/m^2$ $\sigma = 5.67 \times 10^{-8}\ Wm^{-2}K^{-4}$ $\lambda = 0.769$

Line-by-line codes accounting for atmospheric profiles of temperature, water vapor, and aerosols.

Simplified fits to the data, such as for CO_2

$$\boldsymbol{RF = 5.35\, ln\left(\frac{CO_2}{CO_{2\,orig}}\right) W/m^2}$$

Logarithmic because some particular lines are already saturated.

Forcings for lower concentration gases (such as CFCs) are linear in concentration.

RF for a doubling of CO_2 is likely $\boldsymbol{3.7 \pm 0.4\, \frac{W}{m^2}}$

Same order of magnitude as 2% increase of solar forcing

Total forcing from trace greenhouse gases mentioned in Step 3, is currently about $\mathbf{2.5\ W/m^2}$

Net forcing (including cooling impacts of aerosols and natural changes) is $\mathbf{1.6 \pm 1.0\ W/m^2}$ since pre-industrial.

- uncertainty mostly related to aerosol effects.

Current forcings growth dominated by increasing CO_2, with potentially a small role for

- decreases in reflective aerosols (sulphates, particularly in the US and EU)
- increases in absorbing aerosols (like soot, particularly from India and China and from biomass burning)

Step 3 - "CO_2 is up more than 30%, CH_4 has more than doubled, N_2O is up 15%, tropospheric O_3 has also increased. New compounds such as halocarbons (CFCs, HFCs) did not exist in the pre-industrial atmosphere."

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

The CO_2 Answer in 6 Easy Steps

4. **Radiative forcing is a useful diagnostic and can easily be calculated**



The CO_2 Answer in 6 Easy Steps

5. **Climate sensitivity is around 3 C° for a doubling of CO_2**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Climate sensitivity is response of global mean temperature to forcing, $\frac{C°}{W/m^2}$

after 'fast feedbacks' have occurred (atmospheric temperatures, clouds, water vapor, winds, snow, sea ice etc.)

before 'slow' feedbacks have started (ice sheets, vegetation, carbon cycle etc.)

Sensitivity can be assessed from any particular period in the past where

changes in forcing are known

corresponding equilibrium **temperature change** can be estimated

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Last glacial period had large forcing, **~7 W/m^2**
from ice sheets, greenhouse gases, dust and vegetation

Large temperature response, **~5 ºC**

implying a sensitivity of about **3 ºC** with error

3.7 W/m^2 x 5 ºC / 7 W/m^2 = 2.6 ºC

Provided link to 2006 estimate of response to volcanoes, the last millennium, remote sensing etc. which was also **3 ºC**.

"Converting the estimate for doubled CO_2 to a more useful factor gives **~0.75 ºC/(W/m^2)**"

3 ºC / 3.7 W/m^2 = 0.81 ºC/(W/m^2)

5 ºC / 7 W/m^2 = 0.71 ºC/(W/m^2)

1.5 ºC / 5 W/m^2 = 0.3 ºC/(W/m^2)

The CO_2 Answer in 6 Easy Steps

5. **Climate sensitivity is around 3 C° for a doubling of CO_2**



The CO_2 Answer in 6 Easy Steps

6. **Radiative forcing x climate sensitivity is a significant number**

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

Current forcings; **1.6 W/m² x 0.75 ºC/(W/m²) = 1.2 ºC** at equilibrium. Oceans take time to warm up, air is only up **0.7ºC** Remaining **0.5 ºC** 'in the pipeline'.

Also estimated using changes in ocean heat content over last decade (about equal to radiative imbalance) of **~0.7 W/m²**, implying that this 'unrealised' forcing will lead to another

$$0.7\ \mathrm{W/m^2} \times 0.75\ ºC/(\mathrm{W/m^2}) \cong 0.5\ ºC$$

Additional forcings in business-as-usual scenarios range roughly from **3 to 7 W/m²**

Additional warming at equilibrium would be **2 to 5 ºC**.

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/

The CO_2 Answer in 6 Easy Steps

1. There is a natural greenhouse effect
2. Trace gases contribute to the natural greenhouse effect
3. The trace greenhouse gases have increased markedly due to human emissions
4. Radiative forcing is a useful diagnostic and can easily be calculated
5. Climate sensitivity is around 3 C° for a doubling of CO_2
6. Radiative forcing x climate sensitivity is a significant number

realclimate.org/index.php/archives/2007/08/the-co2-problem-in-6-easy-steps/